

## DAFTAR RUJUKAN

- Afrianti, L. H., Sukandar, E. Y., Ibrahim, S., Adnyana, I. K., Teknologi, D., & Universitas, P. (2010). Antihyperuricemia [ Studies on 2-Methylester-1-H-Pyrolle-4-Carboxylic Acid Compound in Ethylacetate Extract of Snake Fruit Variety Bongkok as Antioxidant and Anthyperuricemic ] Metode, *XXI*(1), 66–72.
- Amlashi, H. A., Madani, H., Sonboli, A., & Khaghani, S. (2020). Volatile Composition of The Leaves and Calyces Essential Oil of Roselle ( *Hibiscus sabdariffa* L .) From Iran. *Journal Of Essential Oil Bearing Plants*, *4*, 743–755. <https://doi.org/10.1080/0972060X.2020.1832585>
- Berk, Z. (2018). *Food Process Engineering and Technology*. <https://doi.org/10.1016/B978-0-12-812018-7.00013-0>
- Božović, M., Navarra, A., Garzoli, S., Pepi, F., Ragno, R., Božović, M., ... Ragno, R. (2017). Essential oils extraction: a 24-hour steam distillation systematic methodology. *Natural Product Research*, *64*(9)(April), 0. <https://doi.org/10.1080/14786419.2017.1309534>
- Cahyati, S., Kurniasih, Y., & Khery, Y. (2008). Efisiensi Isolasi Minyak Atsiri dari Kulit Jeruk dengan Metode Distilasi Air-Uap Ditinjau dari Perbandingan Bahan Baku dan Pelarut yang Digunakan. *Jurnal Ilmiah Pendidikan Kimia "Hydrogen,"* *4*(2), 103–110.
- Chandra, A., & Proborini, W. D. (2017). Analisa Komposisi Minyak Atsiri Kulit Jeruk Manis Hasil Ekstraksi Metode Microwave Hydrodiffusion and Gravity Dengan Gc-MS. *Jurnal Reka Buana*, *3*(1), 53–58.
- Dahlia, A. A., Lestari, A., & Rahayu, A. (2012). Identifikasi Morfologi dan Parameter Spesifik Simplisia dan Ekstrak Daun Rosella (*Hibiscus sabdariffa* L.) Asal Kab. Enrekang (Sullawesi Selatan). *Journal Ilmiah Asia*, *04*(02), 159–175.
- Damayanti, A., & Fitriana, A. (2012). Pemungutan Minyak Atsiri Mawar (Rose Oil) Dengan Metode Maserasi. *Jurnal Bahan Alam Terbarukan*, *1*(2), 1–8.
- Dewi, L. K., Friatnasary, D. L., Herawati, W., Nurhadianty, V., & Cahyani, C. (2018). Studi Perbandingan Metode Isolasi Ekstraksi Pelarut dan Destilasi Uap Minyak Atsiri Kemangi terhadap Komposisi Senyawa Aktif. *Jurnal Rekayasa Bahan Alam Dan Energi Berkelanjutan*, *2*(1), 13–19.
- Dian, Y., & Siregar, I. (2011). Ekstraksi dan Uji Stabilitas Zat Warna Alami dari Bunga Kembang Sepatu ( *Hibiscus rosa-sinensis* L ) dan Bunga Rosela ( *Hibiscus sabdariffa* L ), *2*(3).
- Edogbanya, Suleiman, Olorunmola, & Oijagbe. (2019). Comparative Study On The Antimicrobial Effects Of Essential Oils From Peels Of Three Citrus Fruits. *MOJ Biology and Medicine*, *4*(2), 49–54.

<https://doi.org/10.15406/mojbm.2019.04.00113>

- Figueiredo, C., Jose G., B., Luis G., P., & Johannes J., C. S. (2008). Factors affecting secondary metabolite production in plants: volatile components and essential oils. *Flavour and Fragrance Journal*, (April), 213–226. <https://doi.org/10.1002/ffj>
- Frassinetti, S., Caltavuturo, L., Cini, M., Croce Della, C. M., & Maserti, B. E. (2011). Antibacterial and antioxidant activity of essential oils from citrus spp. *Journal of Essential Oil Research*, 23(1), 27–31. <https://doi.org/10.1080/10412905.2011.9700427>
- Goswami, P., Padalia, R. C., Verma, R. S., & Chauhan, AmitChanotiya, C. S. (2016). Essential oil composition of *Artemisia stelleriana* Besser from India, 2905(April), 0–6. <https://doi.org/10.1080/10412905.2016.1175978>
- Hidayati, N., & Syahnandiaratri, H. (2018). Analisis Pengaruh Daya Microwave Pada Proses Pengambilan Minyak atsiri Daun Kelor (*Moringa Oleifera*) Dengan Metode Microwave Assisted Extraction (Mae). *Simposium Nasional RAPI XVII*, 124–129.
- Mahadevan, N., & Kamboj, P. (2009). *Hibiscus sabdariffa* Linn . – An overview, 8(1), 77–83.
- Mahfud, T. (2015). Ekstraksi Pewarna Alami Kelopak Bunga Rosella ( *Hibiscus Sabdariffa* ) Pada Pembuatan Minuman Serbuk Instan Rosella. *Jurnal Sains Terapan*, 1(1), 27–33.
- Mungole, A., & Chaturvedi, A. (2011). *Hibiscus sabdariffa* L. A Rich Source of Secondary Metabolites. *International Journal OF Pharmaceutical Sciences Review and Research*, 6(1), 83–87.
- Nurnasari, E., & Khuluq, D. (2017). Potensi Diversifikasi Rosela Herbal ( *Hibiscus Sabdariffa* L.) untuk Pangan dan Kesehatan. *Jurnal Pertanian*, 9(2).
- Sahala, A., & Soegihardjo, C. J. (2012). Uji Aktivitas Antioksidan Dan Penetapan Kadar Fenolat Total Fraksi Air Daun Ketapang (*Terminalia Catappa* L.) Dengan Metode Dpph (2,2-Diphenyl-1-Picrylhydrazyl) Dan Metode Folin-Ciocalteu. *Jurnal Farmasi Sains Dan Komunitas*, 9(2), 91–97. <https://doi.org/10.1088/1751-8113/44/8/085201>
- Slamet, Ulyarti, & Rahmi. (2019). Pengaruh Lama Fermentasi Daun Nilam Menggunakan Ragi Tempe Terhadap Rendemen Dan Mutu Fisik Minyak Nilam (*Pogostemon Cablin Benth.*). *Jurnal Teknologi Hasil Perikanan Indonesia*, 11(01), 19–25.
- Stenis, D. van. (2006). *Flora*. (I. Moeso & Surjowinoto, Eds.). Jakarta: PT. Pradnya paramitha.
- Tao, N., Yueojin, L., & Z., M. (2009). Chemical composition and antimicrobial activities of essential oil from the peel of bingtang sweet orange (*Citrus sinensis* Osbeck). *International Journal of Food Sciences and Technology*, 44(7), 1281–1285.

- Tumbas, V. T., Četković, G. S., Djilas, S. M., Čanadanović-Brunet, J. M., Vulić, J. J., Knéz, Ž., & Škerget, M. (2010). Antioxidant activity of mandarin (*Citrus reticulata*) peel. *Acta Periodica Technologica*, 41(January), 195–203. <https://doi.org/10.2298/APT1041195T>
- Widyasanti, A., Rohdiana, D., & Ekatama, N. (2016). Aktivitas Antioksidan Ekstrak Teh Putih (*Camellia Sinensis*) Dengan Metode Dpph (2,2 Difenil-1-Pikrilhidrazil) Antioxidant Activities of White Tea Extract (*Camellia sinensis*) Using DPPH (2,2 diphenyl-1-picrylhydrazyl) Method. *Fortech*, 1(1), 2016.
- Wiley, J., Figueiredo, A. C., Barroso, J. G., Pedro, L. G., & Scheffer, J. J. C. (2008). Factors affecting secondary metabolite production in plants : volatile components and essential oils. *Flavour and Fragrance Journal*, (May), 213–226. <https://doi.org/10.1002/ffj>
- Yassir, M. (2019). Jenis Tanaman Obat di Kecamatan Semadam Kabupaten Aceh Tenggara. *Journal Sains Dan Aplikasi*, VII(1), 6–12.

