

DAFTAR REFERENSI

- Anam, K. (2015). Isolasi Senyawa Triterpenoid dari Alga Merah (*Eucheuma cottoni*) Menggunakan Kromatografi Lapis Tipis (KLT) dan Analisisnya Menggunakan Spektrofotometer UV-Vis dan FTIR. *Skripsi*, 22–23.
- Carbone, M., Ciavatta, M. L., Manzo, E., Li, X. L., Mollo, E., Mudianta, I. W., Guo, Y. W., & Gavagnin, M. (2019). Amphilectene diterpene isonitriles and formamido derivatives from the hainan nudibranch *Phyllidia coelestis*. *Marine Drugs*, 17(11). 1-14.
- Darmapatni, K. A. G., Basori, A., & Suaniti, N. made. (2016). Pengembangan Metode Gc-Ms Untuk. *Jurnal Biosains Pascasarjana*, 18(3), 255–270.
- Dewoto, H. R. (2007). *Pengembangan Obat Tradisional Indonesia Menjadi Fitofarmaka* *. Majalah Kedokteran Indonesia. 205–211.
- Gerwick, W. H., & Moore, B. S. (2012). Lessons from the past and charting the future of marine natural products drug discovery and chemical biology. *Chemistry and Biology*, 19(1), 85–98.
- Jaisamut, S., Prabpai, S., Tancharoen, C., Yuenyongsawad, S., Hannongbua, S., Kongsaeree, P., & Plubrukarn, A. (2013). Bridged tricyclic sesquiterpenes from the tubercle nudibranch *Phyllidia coelestis* Bergh. *Journal of Natural Products*, 76(11), 2158–2161.
- Made, D. A. N., Parwata, I. M. O. A., & Parthasutema, I. . M. (2015). Analisis Kadar Metamfetamine pada Sampel Darah dengan Metode GC-MS. *Chemistry Laboratory*, 2, 18–29.
- Mudianta, I. W., Martiningsih, N. W., Prasetia, I. N. D., & Nursid, M. (2016). Bioactive terpenoid from the balinese nudibranch *hypsodoris infucata*. *Indonesian Journal of Pharmacy*, 27(2), 104–110.
- Mukhtarini. (2011). Ekstraksi, pemisahan senyawa, dan identifikasi senyawa aktif. *Jurnal of Pharmacy*, V, 361. 5-8.
- Prawanayoni, S. S., & Sudirga, S. K. (2020). Isolasi Dan Identifikasi Senyawa Antijamur Daun Jeringau (*Acorus calamus* Linn.) Sebagai Pengendali Jamur *Athelia rolfsii* Sacc. Penyebab Penyakit Busuk Batang Pada Tanaman Kedelai. *Metamorfosa: Journal of Biological Sciences*, 7(2), 10-15
- (Patil, R.A., Weatherly, C.A., Armstrong, D.W., 2018.

Chiral GasChromatography, in: Chiral Analysis. Elsevier, pp. 468–505.

- Sim, D. C. M., Wayan Mudianta, I., White, A. M., Martiningsih, N. W., Loh, J. J. M., Cheney, K. L., & Garson, M. J. (2018). New sesquiterpenoid isonitriles from three species of phyllid nudibranchs. *Fitoterapia* Sim, D. C. M., Wayan Mudianta, I., White, A. M., Martiningsih, N. W., Loh, J. J. M., Cheney, K. L., & Garson, M. J. (2018). New Sesquiterpenoid Isonitriles from Three Species of Phyllid nudibranchs. *Fitoterapia*, 126, 69–73.
- Stoffels, B. E. M. W., Van Der Meij, S. E. T., Hoeksema, B. W., Van Alphen, J., Van Alen, T., Meyers-Muñoz, M. A., De Voogd, N. J., Tuti, Y., & Van Der Velde, G. (2016). Phylogenetic relationships within the phyllidiidae (Opisthobranchia, Nudibranchia). *ZooKeys*, 2016(605), 1–35.
- Tsuchida, S. (2002). Test and repair of non-volatile commodity and embedded memories. *IEEE International Test Conference (TC)*, 3(May), 1223. 2-7.
- White, A. M., Pierens, G. K., Skinner-adams, T., Andrews, K. T., Bernhardt, P. V., Krenske, E. H., Mollo, E., & Garson, M. J. (2015). Antimalarial Isocyano and Isothiocyanato Sesquiterpenes with Tri- and Bicyclic Skeletons from the Nudibranch *Phyllidia ocellata*. *Natural Products*, 1(Figure 2), 2–7.
- Wu, Q., Chen, W. T., Li, S. W., Ye, J. Y., Huan, X. J., Gavagnin, M., Yao, L. G., Wang, H., Miao, Z. H., Li, X. W., & Guo, Y. W. (2019). Cytotoxic nitrogenous terpenoids from two South China sea nudibranchs *phyllidiella pustulosa*, *Phyllidia coelestis*, and their sponge-prey *Acanthella cavernosa*. *Marine Drugs*, 17(1), 23–27.