

DAFTAR PUSTAKA

- Achhab, Y. El *et al.* (2019) ‘Prevalence, control and risk factors related to hypertension among moroccan adults: A multicentre study’, *Eastern Mediterranean Health Journal*, 25(7), pp. 447–456. doi: 10.26719/emhj.18.057.
- Angkawijaya, A. A., Pangemanan, J. M. and Siagian, I. E. . (2016) ‘Hubungan Tingkat Pengetahuan Masyarakat dengan Tindakan Pencegahan Hipertensi Di Desa Motoboi Kecil Kecamatan Kotamobagu Selatan PENDAHULUAN Hipertensi atau tekanan darah tinggi merupakan faktor pemicu terjadinya komplikasi seperti stroke , gagal jantung’, *Kedokteran Komunitas dan Tropik*, IV.
- Badego, B., Yoseph, A. and Astatkie, A. (2020) ‘Prevalence and risk factors of hypertension among civil servants in Sidama Zone, south Ethiopia’, *PLoS ONE*, 15(6 June 2020), pp. 1–17. doi: 10.1371/journal.pone.0234485.
- Ban, H. J. and Kim, H. S. (2020) ‘Applying the modified health belief model (Hbm) to korean medical tourism’, *International Journal of Environmental Research and Public Health*, 17(10). doi: 10.3390/ijerph17103646.
- Chakma, T. *et al.* (2017) ‘High prevalence of hypertension and its selected risk factors among adult tribal population in Central India’, *Pathogens and Global Health*, 111(7), pp. 343–350. doi: 10.1080/2047724.2017.1396411.
- Chimberengwa, P. T. and Naidoo, M. (2018) ‘Knowledge, attitudes and practices related to hypertension among residents of a disadvantaged rural community in southern Zimbabwe’, *PLoS ONE*, 14(6), pp. 1–16. doi: 10.1371/journal.pone.0215500.
- DeGuire, J. *et al.* (2019) ‘Blood pressure and hypertension’, *Health Reports*, 30(2), pp. 14–21. doi: 10.25318/82-003-x201900200002.
- Desormais, I. *et al.* (2019) ‘The prevalence, awareness, management and control of

- hypertension in men and women in Benin, West Africa: The TAHES study', *BMC Cardiovascular Disorders*, 19(1), pp. 1–12. doi: 10.1186/s12872-019-01273-7.
- E.Drawz, P. et al. (2020) 'Blood Pressure Measurement: A KDOQI Perspective', *Physiology & behavior*, 63(8), pp. 1–18. doi: 10.1053/j.ajkd.2019.08.030.
- Erkoc, S. B. et al. (2012) 'Hypertension knowledge-level scale (HK-LS): A study on development, validity and reliability', *International Journal of Environmental Research and Public Health*, 9(3), pp. 1018–1029. doi: 10.3390/ijerph9031018.
- Ernawati, I., Fandinata, S. S. and Permatasari, S. N. (2020) 'Translation and validation of the indonesian version of the hypertension knowledge-level scale', *Open Access Macedonian Journal of Medical Sciences*, 8, pp. 630–637. doi: 10.3889/OAMJMS.2020.5152.
- Eva Rayanti, R., Nugroho, K. P. A. and Marwa, S. L. (2021) 'Health Belief Model dan Management Hipertensi Pada Penderita Hipertensi Primer di Papua', *Jurnal Keperawatan Muhammadiyah*, 6(1), p. 51.
- Fiuza, M. (2019) 'Control of risk factors in hypertensive patients: A task to fulfill', *Revista Portuguesa de Cardiologia*, 38(6), pp. 439–440. doi: 10.1016/j.repc.2019.07.004.
- Gupta, R. and Xavier, D. (2018) 'Hypertension: The most important non communicable disease risk factor in India', *Indian Heart Journal*, 70(4), pp. 565–572. doi: 10.1016/j.ihj.2018.02.003.
- Hurst, C. P., Rakkapao, N. and Hay, K. (2020) 'Impact of diabetes self-management, diabetes management self-efficacy and diabetes knowledge on glycemic control in people with Type 2 Diabetes (T2D): A multicenter study in Thailand', *PLoS ONE*, 15(12 December), pp. 1–14. doi: 10.1371/journal.pone.0244692.
- Iyalomhe, G. B. S. and Iyalomhe, S. I. (2010) 'Hypertension-related knowledge, attitudes and life-style practices among hypertensive patients in a sub-urban Nigerian community', *Journal of Public Health and Epidemiology*, 2(4), pp. 71–77.

- Available at: <http://www.academicjournals.org/jphe>.
- Kjellgren, K. I. *et al.* (1997) ‘Hypertensive patients’ knowledge of high blood pressure’, *Scandinavian Journal of Primary Health Care*, 15(4), pp. 188–192. doi: 10.3109/02813439709035026.
- Kosasih, A. *et al.* (2019) *Konsensus Penatalaksanaan Hipertensi 2019*. 2019th edn. Edited by A. Anna Lukito, E. Harmeyaty, and N. M. Hustrini. Jakarta: Indonesian Society of Hypertension. Available at: http://www.inash.or.id/upload/event/event_Update_konsensus_2019123191.pdf.
- Leung, A. A. *et al.* (2019) ‘Risk factors for hypertension in Canada’, *Health Reports*, 30(2), pp. 1–13.
- Limbong, V. A., Rumayar, A. and Kandou, grace D. (2018) ‘Hubungan Pengetahuan Dan Sikap Dengan Kejadian Hipertensi Di Wilayah Kerja Puskesmas Tateli Kabupaten Minahasa’, *Kesmas*, 7(4).
- Malahayati, K., Supriyadi, R. and Sastramihardja, H. S. (2016) ‘Knowledge and Attitude towards Prevention and Management of Hypertension in Jatinangor Sub-district’, *Althea Medical Journal*, 3(2), pp. 222–229. doi: 10.15850/amj.v3n2.801.
- Mirzaei, Mohsen *et al.* (2020) ‘Awareness, treatment, and control of hypertension and related factors in adult Iranian population’, *BMC Public Health*, 20(1), pp. 1–10. doi: 10.1186/s12889-020-08831-1.
- Olack, B. *et al.* (2015) ‘Risk factors of hypertension among adults aged 35–64 years living in an urban slum Nairobi, Kenya’, *BMC Public Health*, 15(1), pp. 1–9. doi: 10.1186/s12889-015-2610-8.
- Oliveria, S. A. *et al.* (2005) ‘Hypertension knowledge, awareness, and attitudes in a hypertensive population’, *Journal of General Internal Medicine*, 20(3), pp. 219–225. doi: 10.1111/j.1525-1497.2005.30353.x.
- Rachman, R. A., Noviati, E. and Kurniawan, R. (2021) ‘Efektifitas Edukasi Health

- Belief Models Dalam Perubahan Perilaku Pasien Hipertensi: Literatur Review’, *Healthcare Nursing Journal*, 3(1), pp. 71–80. doi: 10.35568/healthcare.v3i1.1091.
- Rahman, M. et al. (2018) ‘Prevalence, Treatment Patterns and Risk Factors of Hypertension and Pre-hypertension among Bangladeshi Adults’, 32(5), pp. 334–348. doi: 10.1038/s41371-017-0018-x.Prevalence.
- Saka, M., Shabu, S. and Shabila, N. (2020) ‘Prevalence of hypertension and associated risk factors in older adults in Kurdistan, Iraq’, *Eastern Mediterranean Health Journal*, 26(3), pp. 265–272. doi: 10.26719/emhj.19.029.
- Sayed Fazel, Z. M. et al. (2015) ‘Knowledge, treatment, control, and risk factors for hypertension among adults in Southern Iran’, *International Journal of Hypertension*, 2015.
- Setianingsih, R., Tamtomo, D. and Suryani, N. (2016) ‘Health Belief Model: Determinants of Hypertension Prevention Behavior in Adults at Community Health Center, Sukoharjo, Central Java’, *Journal of Health Promotion and Behavior*, 01(03), pp. 160–170. doi: 10.26911/thejhp.2016.01.03.03.
- Suaib, M., Cheristina and Dewiyanti (2019) ‘Hubungan Tingkat Pengetahuan Dengan Kejadian Hipertensi Pada Lansia’, *Jurnal Fenomena Kesehatan*, 2(1), pp. 269–276.
- Tasić, I. et al. (2020) ‘Predictors of cardiovascular events in hypertensive patients with high cardiovascular risk’, *Medicina (Lithuania)*, 56(4), pp. 1–11. doi: 10.3390/medicina56040182.
- Unger, T. et al. (2020) ‘2020 International Society of Hypertension Global Hypertension Practice Guidelines’, *Hypertension*, 75(6), pp. 1334–1357. doi: 10.1161/HYPERTENSIONAHA.120.15026.
- Wang, J. et al. (2017) ‘Prevalence and risk factors of comorbidities among hypertensive patients in China’, *International Journal of Medical Sciences*, 14(3), pp. 201–212. doi: 10.7150/ijms.16974.

Yogiantoro, M. (2014) *Buku Ajar Ilmu Penyakit Dalam*. 6th edn. Edited by S. Setiari et al. Jakarta: Interna Publishing.

